ELSEVIER

Environmental Pollution 129 (2004) I–IV

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

# Contents of Volume 129

**Number 1 2004**

**New Initiative**

1 Preliminary evidence of the role of hydrogen peroxide in the degradation of benzo[a]pyrene by a non-white rot fungus *Fusarium solani*
E. VEIGNIE, C. RAFIN, P. WOISEL, F. CAZIER (France)
*An alternative metabolic pathway was demonstrated.*

**Research Papers**

5 Evaluation of factors influencing root-induced changes of copper fractionation in rhizosphere of a calcareous soil
S. TAO, W.X. LIU, Y.J. CHEN, F.L. XU, R.W. DAWSON, B.G. LI, J. CAO, X.J. WANG, J.Y. HU, J.Y. FANG (PR China)
*Influence of root exudates on copper fractionation appeared to be due to complexation, rather than acidification or alkalisation.*

13 A case study of contaminants on military ranges: Camp Edwards, Massachusetts, USA
J. CLAUSEN, J. ROBB, D. CURRY, N. KORTE (USA)
*Assessment of groundwater and soil contamination at US military ranges can be limited primarily to explosive-related compounds.*

23 Tissue metal levels in Muskrat (*Ondatra zibethica*) collected near the Sudbury (Ontario) ore-smelters; prospects for biomonitoring marsh pollution
G.H. PARKER (Canada)
*Muskrats appear to be useful bioindicators of metal pollution in semi-aquatic environments.*

31 Constraints on the use of $^{137}Cs$ as a time-marker to support CRS and SIT chronologies
J.M. ABRIL (Spain)
*Simple methods based on the identification of $^{137}Cs$ fallout peaks cannot provide a definite support for CRS and SIT chronologies.*

39 Assessing the cause of impacts on benthic organisms near Rouyn-Noranda, Quebec
U. BORGMANN, M. NOWIERSKI, L.C. GRAPENTINE, D.G. DIXON (Canada)
*Metal bioaccumulation and metals in overlying water during toxicity tests help identify the cause of sediment toxicity and reduced benthic biodiversity.*

49 Ostracoda and foraminifera as short-term tracers of environmental changes in very polluted areas: the Odiel Estuary (SW Spain)
F. RUIZ, M.L. GONZÁLEZ-REGALADO, J. BORREGO, M. ABAD, J.G. PENDÓN (Spain)
*In a polluted estuary, the meiofaunal assemblages experience different changes even in adjacent sedimentary environments, closely related to anthropogenic and natural causes.*

63 Rainwater chemistry at the summit and southern flank of the Itatiaia massif, Southeastern Brazil
W.Z. DE MELLO, M.D. DE ALMEIDA (Brazil)
*Rainwater chemistry at the Itatiaia massif in SE Brazil is strongly dominated by S and N inorganic species, and influenced by human activities.*

69 Low molecular weight thiols in arsenic hyperaccumulator *Pteris vittata* upon exposure to arsenic and other trace elements
Y. CAI, J. SU, L.Q. MA (USA)
*Arsenic induces synthesis of low molecular weight thiols in the arsenic hyperaccumulator* Pteris vittata.

79 Sulphur dioxide adsorption in Scots pine canopies exposed to high ammonia emissions near a Cu–Ni smelter in SW Finland
J. DEROME, T. NIEMINEN, A. SAARSALMI (Finland)
*Increased levels of N and S deposition have increased foliar N and S concentrations and needle size in Scots pine.*

89 Chemical coupling between ammonia, acid gases, and fine particles
B.H. BAEK, V.P. ANEJA, Q. TONG (USA)
*Our study of ammonia, acid gases, and fine particles provided a gas-to-particle conversion process that improved prediction of long-range transport of pollutants, air quality, and climate change.*

99 Cellular energy allocation in zebra mussels exposed along a pollution gradient: linking cellular effects to higher levels of biological organization
R. SMOLDERS, L. BERVOETS, W. DE COEN, R. BLUST (Belgium)
*Exposure of zebra mussels along a pollution gradient has adverse effects on the cellular energy allocation, and results can be linked with higher levels of biological organization.*

113 The study of metal contamination in urban soils of Hong Kong using a GIS-based approach
X. LI, S.-L. LEE, S.-C. WONG, W. SHI (Hong Kong), I. THORNTON (UK)
*GIS can be used to identify soil contamination hot-spot areas and to assess potential pollutant sources in an urban community.*

doi: 10.1016/S0269-7491(04)00076-4

125 The effect of heavy metal exposure on egg size, eggshell thickness and the number of spermatozoa in blue tit *Parus caeruleus* eggs
T. DAUWE, E. JANSSENS (Belgium), B. KEMPENAERS (Germany), M. EENS (Belgium)
*Exposure to elevated levels of heavy metals had a negative effect on the number of spermatozoa in the eggs of free-living blue tits (*Parus caeruleus*).*

131 Spatial and temporal trends in surface water and sediment contamination in the Laurentian Great Lakes
C. MARVIN, S. PAINTER, D. WILLIAMS, V. RICHARDSON (Canada), R. ROSSMANN, P. VAN HOOF (USA)
*While contaminant levels in the Great Lakes have declined over the past 20 years, Lake Ontario remains the most contaminated.*

145 Sulphur, nitrogen and carbon content of *Sphagnum capillifolium* and *Pseudevernia furfuracea* exposed in bags in the Naples urban area
S. VINGIANI, P. ADAMO, S. GIORDANO (Italy)
*The lichen* Pseudevernia furfuracea *is more effective than the moss* Sphagnum capillifolium *as S and N pollutants biomonitor.*

159 The need for ammonia abatement with respect to secondary PM reductions in Europe
J.W. ERISMAN, M. SCHAAP (The Netherlands)
*Besides those for $NO_x$ and $SO_2$, ammonia emissions need to be decreased to reduce secondary fine PM effectively.*

**Volume Number 2 2004**

**Research Papers**

165 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 1. Degradation rates using tropical marine microbes
P. MERCURIO, K.A. BURNS, A. NEGRI (Australia)
*Vegetable-derived lubricants were more degradable than mineral oil lubricants.*

175 Testing the ecotoxicology of vegetable versus mineral based lubricating oils: 2. Induction of mixed function oxidase enzymes in barramundi, *Lates calcarifer*, a tropical fish species
P. MERCURIO (USA), K.A. BURNS (Australia), J. CAVANAGH (New Zealand)
*Vegetable-based lubricating oils appear to cause a tropical fish species less stress than mineral oils.*

183 The ecotoxicology of vegetable versus mineral based lubricating oils: 3. Coral fertilization and adult corals
P. MERCURIO, A.P. NEGRI, K.A. BURNS, A.J. HEYWARD (Australia)
*Vegetable-derived lubricants were less toxic to adult corals than their mineral counterparts.*

195 Spatio-temporal behaviour and mass balance of fluorine in forest soils near an aluminium smelting plant: short- and long-term aspects
M. EGLI, S. DÜRRENBERGER, P. FITZE (Switzerland)
*The effects of F can be seen in charges in soil chemistry, vegetation and clay mineralogy.*

209 Distribution and coassociations of trace elements in soft tissue and byssus of *Mytilus galloprovincialis* relative to the surrounding seawater and suspended matter of the southern part of the Korean Peninsula
P. SZEFER, B.-S. KIM (Poland), C.-K. KIM, E.-H. KIM, C.-B. LEE (South Korea)
*Mytilus galloprovincialis can be used as a biomonitor for trace elements in coastal waters off Korea.*

229 Role of manganese oxides in the exposure of mute swans (*Cygnus olor*) to Pb and other elements in the Chesapeake Bay, USA
W.N. BEYER, D. DAY (USA)
*Lead was scavenged by Mn and Fe oxides and deposited on aquatic vegetation.*

237 Overexpression of *LCT1* in tobacco enhances the protective action of calcium against cadmium toxicity
D.M. ANTOSIEWICZ, J. HENNIG (Poland)
*This is the first report of LCT1 in Ca acquisition and regulation of Cd-toxicity by Ca.*

247 The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon
L.D. LACERDA, M. DE SOUZA, M.G. RIBEIRO (Brazil)
*Land use change is a strong re-mobilizing agent of Hg distribution in Amazon soils.*

257 Antimony distribution and environmental mobility at an historic antimony smelter site, New Zealand
N.J. WILSON, D. CRAW, K. HUNTER (New Zealand)
*High levels of antimony in primitive smelter soils remain largely immobile on the metre scale.*

267 Degradation of terbutylazine (2-chloro-4-ethylamino-6-terbutylamino-1,3,5-triazine), deisopropyl atrazine (2-amino-4-chloro-6-ethylamino-1,3,5-triazine), and chlorinated dimethoxy triazine (2-chloro-4,6-dimethoxy-1,3,5-triazine) by zero valent iron and electrochemical reduction
T. DOMBEK, D. DAVIS, J. STINE, D. KLARUP (USA)
*Reductive dechlorination via zero valent iron is dependent upon solution pH and the specific triazine.*

277 Attenuation reactions in a multiple contaminated aquifer in Bitterfeld (Germany)
S. HEIDRICH, H. WEIß, A. KASCHL (Germany)
*Attentuation reactions occur in the subsurface and must be investigated on a site-specific basis.*

289 Patterns of metal soil contamination and changes in terrestrial cryptogamic communities
D. CUNY, F.-O. DENAYER, B. DE FOUCAULT (France), R. SCHUMACKER (Belgium), P. COLEIN, C. VAN HALUWYN (France)
*Bryophyte- and lichen-assemblages provide a bioindicative value of heavy metal polluted soils.*

299 Autometallographic tracing of mercury in frog liver
N.S. LOUMBOURDIS, G. DANSCHER (Greece, Denmark)
*Frogs can be good bioindicators of mercury.*

305 Methane oxidation and formation of EPS in compost: effect of oxygen concentration
J.H. WILSHUSEN, J.P.A. HETTIARATCHI (Canada), A. DE VISSCHER (Belgium), R. SAINT-FORT (Canada)
*1.5% oxygen resulted in a slightly higher and more stable methane oxidation activity.*

315 Tributyltin in environmental samples from the Former Derecktor Shipyard, Coddington Cove, Newport RI
T.L. WADE, S.T. SWEET, J.G. QUINN, R.W. CAIRNS, J.W. KING (USA)
*TBT is likely to continue to be bioavailable for many years.*

321 Single-well reactive tracer test and stable isotope analysis for determination of microbial activity in a fast hydrocarbon-contaminated aquifer
L. BURBERY (UK), G. CASSIANI, G. ANDREOTTI, T. RICCHIUTO (Italy), K.T. SEMPLE (UK)
*Reaction rates calculated by push-pull tests are not uniformly distributed in time and space.*

331 The effects of heavy metal contamination on the soil arthropod community of a shooting range
M. MIGLIORINI, G. PIGINO, N. BIANCHI, F. BERNINI, C. LEONZIO (Italy)
*Significant relationships were found between lead accumulation in soil from a shooting range and inhabiting arthropod communities.*

**Number 3 2004**

**Invited Paper**

341 Heavy metals—geochemical bogey men?
M.E. HODSON (UK)
*Serveral alternatives are presented for poorly defined term "heavy metals".*

**Research Papers**

345 Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms
P.L. KLERKS, J.A. NYMAN, S. BHATTACHARYYA (USA)
*Hydrocarbon measurements cannot be substituted for actual toxicity determinations.*

355 Do laboratory salinity tolerances of freshwater animals correspond with their field salinity?
B.J. KEFFORD, P.J. PAPAS, L. METZELING, D. NUGEGODA (Australia)
*Acute laboratory salinity tolerances relate to maximum salinity where organisms occur in nature.*

363 Earthworm biomass as additional information for risk assessment of heavy metal biomagnification: a case study for dredged sediment-derived soils and polluted floodplain soils
B. VANDECASTEELE, J. SAMYN, P. QUATAERT, B. MUYS, F.M.G. TACK (Belgium)
*Polluted clayey dredged sediment-derived soils have a relatively low risk for heavy metal biomagnification due to slow earthworm colonisation.*

377 Biomarker responses of the earthworm *Aporrectodea tuberculata* to copper and zinc exposure: differences between populations with and without earlier metal exposure
T. LUKKARI, M. TAAVITSAINEN, M. SOIMASUO, A. OIKARI, J. HAIMI (Finland)
*The endpoints were analytically reproducible and sensitive parameters.*

387 Using experimental and forest soils to investigate the uptake of polycyclic aromatic hydrocarbons (PAHs) along an urban-rural gradient
F. WONG, T. HARNER, Q.-T. LIU, M.L. DIAMOND (Canada)
*PAH concentrations decreased with distance from the urban centre.*

399 Coral skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for marine pollution by toxic anti-biofouling paints
M. INOUE, A. SUZUKI, M. NOHARA, H. KAN (Japan), A. EDWARD (Micronesia), H. KAWATA (Japan)
*A new index of coral marine pollution is proposed.*

409 Risk assessment of heavy metal pollution for detritivores in floodplain soils in the Biesbosch, the Netherlands, taking bioavailability into account
P.H.F. HOBBELEN, J.E. KOOLHAAS, C.A.M. VAN GESTEL (The Netherlands)
*Low bioavailability reduced the impact on detritivores.*

421 The intestinal parasite *Pomphorhynchus laevis* (Acanthocephala) from barbel as a bioindicator for metal pollution in the Danube River near Budapest, Hungary
F. THIELEN, S. ZIMMERMANN (Germany), F. BASKA (Hungary), H. TARASCHEWSKI, B. SURES (Germany)
*Acanthocephalans of fish may be useful as sentinels of metal pollution in aquatic systems.*

431 Persistent organochlorine residues in human breast milk from Hanoi and Hochiminh city, Vietnam: contamination, accumulation kinetics and risk assessment for infants
N.H. MINH, M. SOMEYA, T.B. MINH, T. KUNISUE, H. IWATA, M. WATANABE, S. TANABE (Japan), P.H. VIET, B.C. TUYEN (Vietnam)
*It is suggested that daily intake rates of persistent organochlorines in mothers in Vietnam may result in health risk for nursing children.*

443 Biokinetics of cadmium, selenium, and zinc in freshwater alga *Scenedesmus obliquus* under different phosphorus and nitrogen conditions and metal transfer to *Daphnia magna*
R.-Q. YU, W.-X. WANG (Hong Kong)
*Phosphorous enrichment influences metal uptake by algae, but not transfer to a zooplankton grazer.*

457 Effects of storage method and duration on the toxicity of marine sediments to embryos of *Crassostrea gigas* oysters
O. GEFFARD, E. HIS, H. BUDZINSKI, J.F. CHIFFOLEAU, A COYNEL, H. ETCHEBER (France)
*Freezing and freeze-drying increased the toxicity of sediments.*

467 Influences of dissolved and colloidal organic carbon on the uptake of Ag, Cd, and Cr by the marine mussel *Perna viridis*
J.-F. PAN, W.-X. WANG (China)
*Organic carbon influences uptake of metals by marine mussels.*

479 Arsenic mobility in contaminated lake sediments
N.P. NIKOLAIDIS (Greece), G.M. DOBBS, J. CHEN, J.A. LACKOVIC (USA)
*Arsenic accumulates at the ground water/lake interface, where it forms insoluble precipitates.*

489 Source apportionment of ambient non-methane hydrocarbons in Hong Kong: Application of a principal component analysis/absolute principal component scores (PCA/APCS) receptor model
H. GUO, T. WANG, P.K.K. LOUIE (Hong Kong)
*The PCA/APCS model provides an easy approach to identify source profiles and to estimate source strengths using ambient samples.*

499 Use of autotrophic sulfur-oxidizers to remove nitrate from bank filtrate in a permeable reactive barrier system
H.S. MOON, K.-H. AHN, S. LEE, K. NAM, J.Y. KIM (South Korea)
*An in situ biological reactive barrier system treating nitrate-contaminated bank filtrate is evaluated.*

I Contents of Volume 129